

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education Ordinary Level

CANDIDATE NAME		
CENTRE NUMBER	CANDIDATE NUMBER	



CHEMISTRY

5070/02

Paper 2 Theory

October/November 2008

1 hour 30 minutes

Candidates answer on the Question Paper.

Additional Materials: Answer Booklet/Paper

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in. Write in dark blue or black pen.

You may use a soft pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Section A

Answer all questions.

Write your answers in the spaces provided in the Question Paper.

Section B

Answer any three questions.

Write your answers on any lined pages and/or separate answer paper.

A copy of the Periodic Table is printed on page 20.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

For Exam	iner's Use
Section A	
В7	
В8	
В9	
B10	
Total	

This document consists of 20 printed pages.



Section A

For Examiner's Use

Answer **all** the questions in this section in the spaces provided.

The total mark for this section is 45.

A1 The diagram shows part of the Periodic Table.

										Не
					В	С	N	0	F	Ne
					Αl	Si	Р	S	Cı	Ar
Fe	Со	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
									I	Xe

Answer these questions using **only** the elements shown in the diagram.


Each element can be used once, more than once or not at all.

Write the symbol for	W	/rite	the	svm	bol	for
----------------------	---	-------	-----	-----	-----	-----

(i)	an element which is in Group 5 and Period 3,		[1]
(ii)	an element which is used as a gas in balloons,		[1]
(iii)	an element which forms ions in aqueous solution which give a white precipitate on reaction with aqueous silver nitrate,		[1]
(iv)	an element which forms an ion of type X^{3-} ,		[1]
(v)	an element which is a catalyst for the hydrogenation of alkenes,		[1]
(vi)	two elements which combine to form a compound which causes acid rain.		[1]
		[Tota	l· 61

A2 Several small pieces of magnesium are placed on a block of solid carbon dioxide. The solid carbon dioxide is at a temperature of -60 °C. The magnesium is ignited and another block of solid carbon dioxide is immediately placed on top.

For Examiner's Use



A vigorous reaction is observed.

dioxide.

$$2Mg + CO_2 \rightarrow 2MgO + C$$

(a) Suggest what could be seen as the reaction proceeds to completion.

[2]

(b) Why is another block of solid carbon dioxide placed above the burning magnesium?

[1]

(c) State one factor in the experiment which slows down the reaction.

[1]

(d) When 2 moles of magnesium react with one mole of carbon dioxide, 810 kJ of energy are released.

Calculate the energy released when 2.0 g of magnesium reacts completely with carbon

[2]

(e)	In a second experiment 6.0 g of magnesium and 4.4 g of carbon dioxide are used. Which solid, magnesium or carbon dioxide is in excess? Show your working.	For Examiner's Use
	[2]	
(f)	Explain, in terms of the energy changes taking place in both bond-making and bond-breaking, why the reaction is exothermic.	
	[2]	
	[Total: 10]	

© UCLES 2008

А3		lousehold waste can be disposed of by being dumped into landfill sites, recycled or burnt. a landfill site, bacteria break down vegetable waste to produce a mixture of gases. Exami Us						
	(a)	Nam	e two gases which are likely to be formed by this bacterial action.					
			and [2]					
	(b)		nall amount of butanoic acid is also formed by bacterial action in landfill sites. Draw tructure of butanoic acid.					
			[1]					
	(c)	at 35	be of 'oil' can be made from the cellulose in waste paper. The waste paper is heated 50 °C under high pressure and in the presence of a nickel catalyst. The equation for reaction is shown.					
			$6C_6H_{10}O_5 \rightarrow C_{22}H_{22}O_2 + 19H_2O + CO_2 + 7CO + 6C$ 'oil'					
		(i)	State the function of a catalyst.					
			[1]					
			The 'oil', $C_{22}H_{22}O_2$, can be used for heating. Write an equation for the complete combustion of this 'oil'.					
			[2]					
			[Total: 6]					

A4 The head of a safety match contains potassium chlorate and antimony sulphide. The side of the matchbox contains red phosphorus.

When a match is struck on the side of the box, the friction produces enough heat to light the match.

For Examiner's Use

(a) The equation for this reaction is show	(a)	The	equation	for this	reaction	is	show
--	-----	-----	----------	----------	----------	----	------

$$5 \text{KC} l \text{O}_3 + 6 \text{P} \rightarrow 5 \text{KC} l + 3 \text{P}_2 \text{O}_5$$
 potassium chlorate phosphorus(V) oxide

Which is the oxidant and which is the reductant in this reaction? Explain your answer.

oxidant	
reductant	
explanation	
	.[2]

- **(b)** Phosphorus(V) oxide, P_2O_5 , absorbs water from the air to form meta-phosphoric acid, HPO₃.
 - (i) Write an equation for this reaction.

[1]


(ii) On addition of more water, phosphoric acid is formed. Phosphoric acid has typical acidic properties. What would you observe when aqueous phosphoric acid is added to

aqueous sodium carbonate,

blue litmus paper?

[2]

(c) Part of the chain structure of antimony sulphide is shown below.



For Examiner's Use

Deduce the empirical formula of antimony sulphide.

.....[1]

[Total: 6]

A5 Cement is made by heating clay with crushed calcium carbonate. During this process, the calcium carbonate is first converted to calcium oxide.

For Examiner's Use

$$CaCO_3 \rightarrow CaO + CO_2$$

(a) (i) What name is given to this type of chemical reaction?

[1]

(ii) Suggest why calcium oxide is used to neutralise acidic soils.

.....[1]


- **(b)** Concrete is made from cement, sand and water. When set, concrete is slightly porous. When rain water soaks through concrete, some of the uncombined calcium oxide dissolves to form calcium hydroxide.
 - (i) Write an equation for this reaction.

[1]

(ii) The aqueous calcium hydroxide in wet concrete reacts with carbon dioxide in the air.

$$Ca(OH)_2 + CO_2 \rightarrow CaCO_3 + H_2O$$

The diagram shows the pH at various points inside a cracked concrete beam.



Describe and explain the change in pH from the surface to the centre of the beam.

.....

্রি

(iii)	25.0 cm3 of an aqueous solution of calcium hydroxide is exactly neutralised by
	18.0 cm ³ of 0.040 mol/dm ³ hydrochloric acid.

For Examiner's Use

$$Ca(OH)_2 + 2HCl \rightarrow CaCl_2 + 2H_2O$$

Calculate the concentration, in mol/dm³, of the aqueous calcium hydroxide.

concentration =mol/dm³ [3]

[Total: 9]

A6	Electrolysis is used to produce many important chemicals such as chlorine, sodium hydroxide and aluminium.						
	(a)	Chlo	orine is used in both water treatment and as a bleach.				
		(i)	Why is chlorine used in water treatment?				
		<i>(</i>)	[1]				
		(ii)	Name a substance, other than chlorine, that is used to bleach wood pulp.				
			[1]				
	(b)	Chlo	orine is used to make chloroethene.				
			H Cl				
			$\overset{\circ}{\mathbf{C}} = \overset{\circ}{\mathbf{C}}$				
			н н				
			proethene can be polymerised to form poly(chloroethene). w a section of a poly(chloroethene) chain to show at least two repeating units.				
			[1]				
	(c)	In th	ne production of aluminium, sodium hydroxide is used to separate aluminium oxide				
	(0)	from	n the impurities in the bauxite ore. The main impurity in the ore is iron(III) oxide.				
		Sug	minium oxide is an amphoteric oxide whilst iron(III) oxide is a basic oxide. gest how these two oxides can be separated by the addition of aqueous sodium roxide.				
			[2]				
			[-]				

(a)	cryolite. What is the function of the cryolite?	For Examiner's Use
	[1]	
(e)	Acidic foods can be safely packed in aluminium containers. Explain why the acid in the food does not attack the aluminium, despite the fact that aluminium is a reactive metal.	
	[2]	
	[Total: 8]	

Section B

Answer three questions from this section.

The total mark for this section is 30.

B7 Ammonia is made by the Haber process using an iron catalyst.

$$N_2 + 3H_2 \rightleftharpoons 2NH_3 \quad \Delta H = -92 \text{ kJ/mol}$$

- (a) On the same axes draw energy profile diagrams to show both the catalysed and the uncatalysed reaction. Label the diagram to show
 - the catalysed and uncatalysed reactions,
 - · the reactants and products,
 - the enthalpy change for the reaction.

[3]

[1]

[2]

- **(b)** The raw materials for the Haber process can be obtained from the air and from hydrocarbons produced by the distillation of petroleum.
 - (i) Describe how pure nitrogen can be separated from other gases in the air.
 - (ii) Describe how hydrogen can be made from hydrocarbons.
- (c) Explain how the position of equilibrium in the Haber process is altered by
 - (i) an increase in pressure, [2]
 - (ii) an increase in temperature. [2]

[Total: 10]

B8	Sorrel	is a	small	areen	plant.
	COLLCI	io a	Jilian	groon	piai it.

- (a) The pigments in the sorrel leaf can be separated by chromatography.
 - (i) Describe how chromatography can be used to separate different pigments. [2]
 - (ii) Explain what is meant by $R_{\rm f}$ value. [1]
- (b) Sorrel plants contain a poisonous carboxylic acid X.
 What can be deduced about X from each of the following three pieces of information?
 - (i) When **X** is warmed with acidified potassium manganate(VII), the solution changes from pink to colourless. [1]
 - (ii) Aqueous bromine is not decolourised when added to a solution of **X**. [1]
 - (iii) A 0.1 mol/dm³ solution of **X** has a pH of 3 whereas a 0.1 mol/dm³ solution of hydrochloric acid has a pH of 1. [1]
- (c) Analysis of 10.0 g of carboxylic acid **X** shows that it contains 2.67 g carbon, 0.220 g hydrogen and 7.11 g oxygen.
 - (i) Deduce the empirical formula of **X**. [3]
 - (ii) The relative molecular mass of **X** is 90. Deduce the molecular formula of **X**. [1]

[Total: 10]


© UCLES 2008 5070/02/O/N/08 **[Turn over**

B9 Electrolysis can be used to remove unwanted hair. The customer holds a metal bar which acts as a positive electrode. A needle, which acts as the negative electrode, is held by the operator.



- (a) What do you understand by the term *electrolysis*?
- (b) The solution around the tip of the needle is mainly a dilute aqueous solution of sodium chloride.

[1]



© UCLES 2008 5070/02/O/N/08

- (i) Name all the ions present in the solution during this electrolysis. [1]
- (ii) During electrolysis a small amount of chlorine is formed at the surface of the skin. Write an ionic equation for this reaction. [1]
- (iii) During electrolysis, a gas forms at the tip of the needle and the solution changes from pH 7 to pH 10.Explain both these changes.[2]
- (c) Explain why aqueous sodium chloride solution conducts electricity but solid sodium chloride does not. [2]
- (d) The sweat glands in the skin produce small amounts of lactic acid.



lactic acid

Lactic acid reacts with ethanol to form an ester.

- (i) State the conditions needed to form an ester. [2]
- (ii) Draw the structure of the ester produced by the reaction of lactic acid with ethanol. [1]

[Total: 10]

B10 Radioactive iodine is used to treat some cancerous tumours.

elements.

(a)	Two radioactive isotopes of iodine are $^{125}_{53}I$ and $^{131}_{53}I$.	
	For each isotope state the type and number of subatomic particles present.	[2]
(b)	Name a reagent that reacts with iodide ions to form iodine molecules. Describe the colour change that occurs in this reaction.	[2]
(c)	Zinc can reduce iodine to iodide ions. Write an ionic equation for this reaction.	[2]
(d)	In cancer treatment, the radioactive iodine can be injected into the tumour with a titan needle.	ium

- (i) Titanium is a transition element. State three characteristic properties of transition
- (ii) An oxide of titanium is formed from Ti³⁺ ions and oxide ions.

 Deduce the formula of this compound. [1]
- (iii) Titanium(IV) chloride, $TiCl_4$, reacts with water to form titanium(IV) oxide, TiO_2 , and hydrogen chloride. Write an equation for this reaction. [1]

[Total: 10]

[2]

© UCLES 2008 5070/02/O/N/08

If you use these lined pages to complete an answer to any question, the question number must be clearly shown.	For Examiner's Use

For
Examiner's Use
1

For
Examiner's Use

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of he Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

ū DATA SHEET

									1 1	
		0	4 He lium	20 Ne Neon	40 Ar Argon	84 Kr Krypton 36	131 Xe Xenon 54	222 Ra Radon 86		175 Lu
		II/		19 T Fluorine	35.5 C1 Chlorine	80 Br Bromine	127 I lodine 53	210 At Astatine 85		Yb
		I		16 Oxygen	32 Sulphur	Selenium	128 Te Tellurium 52	209 Po Polonium 84		169 T H
		>		14 Nitrogen 7	31 P Phosphorus 15	75 AS Arsenic 33	122 Sb Antimony 51	209 Bi Bismuth		167 Er
		2		12 Carbon	28 Si Silicon	73 Ge Germanium	Sn Tin 50	207 Pb Lead		165 Ho
		Ш		11 Boron	27 A1 Aluminium 13	70 Ga Gallium 31	115 In Indium 49	204 T 1 Thallium 81		162 Dy
ts						65 Zn Zinc 30	Cd Cadmium 48	201 Hg Mercury 80		159 Tb
The Periodic Table of the Elements						64 Copper 29	108 Ag Silver	197 Au Gold		157 Gd
e of the	Group					59 Ni Nickel	106 Pd Palladium 46	195 Pt Patinum 78		152 Eu
dic Tabl	Gre					59 Co Cobalt	103 Rh Rhodium 45	192 Ir Iridium		Sm
he Perio			1 T Hydrogen			56 Fe Iron 26	Ru Ruthenium 44	190 Os Osmium 76		147 Pm
T						55 Wn Manganese 25	Tc Technetium 43	186 Re Rhenium		4 B
						52 Cr Chromium 24	96 Mo Molybdenum 42	184 W Tungsten 74		141 P.
						51 V Vanadium 23	93 Nb Niobium 41	181 Ta Tantalum		140 Q
						48 Ti	91 Zr Zirconium 40	178 #f Hafnium 72		
						45 Sc Scandium 21	89 × Yttrium 39	139 La Lanthanum 57	227 AC Actinium 89	d series series
		=		9 Be Berylium 4	24 Mg Magnesium 12	40 Ca Calcium 20	Strontium	137 Ba Barium 56	226 Ra Radium 88	* 58-71 Lanthanoid series † 90-103 Actinoid series
		-		7 Li Lithium	23 Na Sodium	39 K Potassium 19	Rb Rubidium 37	133 Cs Caesium 55	223 Fr Francium 87	* 58–71 † 90–103
	0 00	00				E070/00	/O/NI/00			

58-/1 Lanthanoid series	† 90–103 Actinoid series	
k	_	

175 **Lu** Lutetium

169 Thulium

167 **Er** Erbium

258

Md

Mendelevium
101

257 **Fm** Fermium 100

251 Californium 98

			28
	๙	a = rela ive atomic mass	
Key	×	X = atomic symbol	
	q	b = atomic (proton) number	06

	Cerium 58	Praseodymium 59	Neodymium 60	Promethium 61	Samarium 62	Europium 63	Gadolinium 64	9
a = rela ive atomic mass	232	231	238	237	244	243	247	
X = atomic symbol	┖	Ъа	-	d N	Pu	Am	CB	
and described to the second	Thorium	Protactinium	Uranium	Neptunium	Plutonium	Americium	Curium	
b = atomic (proton) number	06	91	92	93	94	92	96	σ

The volume of one mole of any gas is 24dm3 at room temperature and pressure (r.t.p.).

© UCLES 2008

5070/02/O/N/08